

THE MAGIC (PSILOCYBIN) MUSHROOM INTERACTION REPORT

Created and maintained by a team of licensed pharmacists with PharmD degrees, residency training and board certifications. All hold expertise in psilocybin pharmacology and drug interactions.

[Med]:

The risk of using this medication in combination with psilocybin is expected to be low to moderate. This type of medication has been shown in research to reduce the effects or experience of MACRO-dosed psilocybin for many people. However, some do not experience reduced psilocybin effects while taking this medication. The risk of using this medication with MICRO-dosed psilocybin has not been well studied, but is expected to be low. Contrary to widespread belief, the risk of serotonin syndrome with this medication and psilocybin is low. However, if you are taking other serotonin-increasing medications, the overall risk is higher.

[Med]:

The risk of using this medication with MACRO-dosed psilocybin is expected to be high. This combination should generally be avoided. This medication may lead to accidents and injuries during a psilocybin journey. The medication may abort or severely reduce the journey's effects. In rare/extreme emergencies, this medication might be used to abort a journey, but doing this alone can be very risky. The risk of using this medication with MICRO-dosed psilocybin is expected to be low.

[Med]:

The risk of using this medication with MACRO-dosed psilocybin is expected to be moderate to high since the medication may abort or severely reduce the effects of psilocybin. This combination should generally be avoided. The risk of using this medication with MICRO-dosed psilocybin is unknown, but not expected to cause toxicity. Psilocybin can increase risks (and thus is not recommended) with certain behavioral health conditions that this medication treats.

[Med]:

The risk of using this medication with psilocybin is expected to be low. Based on available evidence and based on what is known about how this medication and psilocybin work, important/serious interactions are not anticipated.

[Med]:

The risk of using this substance with MACRO-dosed psilocybin is expected to be moderate to high. The support of a guide/facilitator is advisable with this combination. This substance may intensify a MACRO-dosed psilocybin journey, or increase the risk of an uncomfortable experience or psychotic symptoms during a journey. This substance and MACRO-dosed psilocybin may increase the chances of 'flashbacks' to the psilocybin experience. The interaction risk of this substance in combination with MICRO-dosed psilocybin is expected to be low.

COPYRIGHT NOTICE: All rights reserved. Any unauthorized use or reproduction of this report or its contents may result in legal action. Neither this report nor its contents, in whole or in part, may be copied, reproduced, distributed, or published in any format or by any means, including but not limited to photocopying, recording, or other electronic or mechanical methods, without the prior written permission of the copyright holder, except in the case of noncommercial uses permitted by copyright law. **You may share this report for the purposes of coordinating or informing harm reduction or healthcare activities related to the use of psilocybin or psilocybin mushrooms by the individual for which this report was intended.**